

# Recursion relations for Hylleraas three-electron integral

Krzysztof Pachucki\* and Mariusz Puchalski†

*Institute of Theoretical Physics, Warsaw University, Hoża 69, 00-681 Warsaw, Poland*

Ettore Remiddi‡

*Dipartimento di Fisica, Università di Bologna,  
and INFN Sezione di Bologna, I-40126 Bologna, Italy  
CERN-PH-TH, CH-1211 Geneva 23, Switzerland*

## Abstract

Recursion relations for Hylleraas three-electron integral are obtained in a closed form by using integration by parts identities. Numerically fast and well stable algorithm for the calculation of the integral with high powers of inter-electronic coordinates is presented.

PACS numbers: 31.25.-v, 31.15.Pf, 02.70.-c

---

\*krp@fuw.edu.pl

†mpuchals@fuw.edu.pl

‡remiddi@bo.infn.it

## I. INTRODUCTION

The explicitly correlated Hylleraas basis set [1] is one of the most efficient representation of a few-electron wave function. It has been applied mostly to helium [2] and lithium atoms [3]. While the general analytic formula for two-electron Hylleraas integrals is quite simple [4], the three-electron case is much more complicated. Fromm and Hill in [5] were able to derive a closed form formula for the generic integral

$$\int d^3 r_1 \int d^3 r_2 \int d^3 r_3 e^{-w_1 r_1 - w_2 r_2 - w_3 r_3 - u_1 r_{23} - u_2 r_{31} - u_3 r_{12}} r_{23}^{-1} r_{31}^{-1} r_{12}^{-1} r_1^{-1} r_2^{-1} r_3^{-1}, \quad (1)$$

which consists of the sum of many multi-valued dilogarithmic functions. This formula was later simplified by Harris [6] to a more symmetric form, in which many spurious singularities have been eliminated. However, only few preliminary results have been reported so far with these functions, namely in Refs. [7, 8]. The more convenient basis set for applications to 3- or 4-electron atoms happen to consist of functions with all  $u_i$  equal to 0, and include polynomials in  $r_{ij}$  and additionally in  $r_i$ , namely the 3-electron Hylleraas functions

$$\phi_{\{n\}} = e^{-w_1 r_1 - w_2 r_2 - w_3 r_3} r_{23}^{n_1-1} r_{31}^{n_2-1} r_{12}^{n_3-1} r_1^{n_4-1} r_2^{n_5-1} r_3^{n_6-1}. \quad (2)$$

A compact expression for the integral with all  $n_i = 0$  and with  $n_1 = 2, n_2 = 0, n_3 = 2, n_4 = n_5 = n_6 = 1$  was derived in [9]. No attempts have been reported so far to derive analytic formulas for larger values of  $n_i$ . They can in principle be obtained from the general formula by Fromm and Hill in Ref. [5] by differentiation with respect to  $u_i$  and  $w_i$ . In practice that is quite complicated for large powers of  $r_{ij}$ . For testing results obtained in this work we have written a Mathematica code for derivation of Hylleraas type of integrals using the Fromm-Hill formula, and obtained results for all  $n_i$  such that  $n_1 + n_2 + n_3 + n_4 + n_5 + n_6 < 10$ . Despite the existence of an analytical result for the generic integral (1), the most advanced results for lithium have been obtained so far by Yan and Drake in series of papers [3], where Hylleraas integrals were calculated by series expansion in angular momenta.

For helium atom recursion relations [4] are available, which allow to express Hylleraas integrals with  $n_1 + n_2 + n_3 = N$  [see Eq. (A5)] as a linear combination of integrals from the lower shell  $N - 1$ . In this work we derive a similar complete set of recursion relations for the three-electron Hylleraas integrals

$$f(n_1, n_2, n_3; n_4, n_5, n_6) = \int \frac{d^3 r_1}{4\pi} \int \frac{d^3 r_2}{4\pi} \int \frac{d^3 r_3}{4\pi} e^{-w_1 r_1 - w_2 r_2 - w_3 r_3} r_{23}^{n_1-1} r_{31}^{n_2-1} r_{12}^{n_3-1} r_1^{n_4-1} r_2^{n_5-1} r_3^{n_6-1}. \quad (3)$$

These recursions allow to eliminate the principal drawback of Hylleraas basis sets, namely the complexity of various integrals for nonrelativistic matrix elements for the lithium atom. Moreover they can be applied also to more than three electron atoms with the condition that the basis set contains at most one odd power of  $r_{ij}$ . In the next section we present our main result, a complete set of recursion relations for the three-electron Hylleraas integrals defined in Eq. (3). In Section III we analyze boundary terms and two-electron recursions, and in Section IV we sketch the derivation of the main result. Finally in Section V we analyze numerical stability and present numerical results for some selected integrals.

## II. THREE-ELECTRON RECURSION RELATIONS

The recursion relations are conveniently divided into two sets. The first set of three recursions increases by 2 any of the first three indices  $n_1, n_2, n_3$  at  $n_4 = n_5 = n_6 = 0$ , starting from the shell  $n_1 + n_2 + n_3 = 0$  (as initial conditions, the values with  $n_i = 0, 1, i = 1, 2, 3$  and  $n_4 = n_5 = n_6 = 0$  are therefore needed); with the notation of Appendix A, they read

$$\begin{aligned}
f(n_1, n_2, n_3 + 2; 0, 0, 0) = & \frac{1 + n_3}{2} \left\{ \right. \\
& \frac{1}{w_1^2} \left[ \frac{n_2(n_2 - 1)}{n_3 + 1} f(n_1, n_2 - 2, n_3 + 2; 0, 0, 0) \right. \\
& + (n_1 + 2n_2 + n_3 + 2) f(n_1, n_2, n_3; 0, 0, 0) \\
& + \frac{1}{n_3 + 1} f(n_1, n_2, n_3 + 2; \star, 0, 0) \\
& + \frac{n_2(n_2 - 1)}{n_1 + 1} f(n_1 + 2, n_2 - 2, n_3; 0, 0, 0) \\
& + \frac{1}{n_1 + 1} f(n_1 + 2, n_2, n_3; 0, 0, \star) \\
& - \frac{\delta_{n_2}}{n_3 + 1} f(n_1, \star, n_3 + 2; 0, 0, 0) \\
& \left. - \frac{\delta_{n_2}}{n_1 + 1} f(n_1 + 2, \star, n_3; 0, 0, 0) \right] \\
& + \frac{1}{w_2^2} \left[ \frac{n_1(n_1 - 1)}{n_3 + 1} f(n_1 - 2, n_2, n_3 + 2; 0, 0, 0) \right. \\
& + (2n_1 + n_2 + n_3 + 2) f(n_1, n_2, n_3; 0, 0, 0) \\
& + \frac{1}{n_3 + 1} f(n_1, n_2, n_3 + 2; 0, \star, 0) + \\
& \left. \frac{n_1(n_1 - 1)}{n_2 + 1} f(n_1 - 2, n_2 + 2, n_3; 0, 0, 0) + \right.
\end{aligned}$$

$$\begin{aligned}
& \frac{1}{n_2 + 1} f(n_1, n_2 + 2, n_3; 0, 0, \star) \\
& - \frac{\delta_{n_1}}{n_3 + 1} f(\star, n_2, n_3 + 2; 0, 0, 0) \\
& - \frac{\delta_{n_1}}{n_2 + 1} f(\star, n_2 + 2, n_3; 0, 0, 0) \Big] \\
& - \frac{w_3^2}{w_1^2 w_2^2} \Big[ \frac{n_3 (n_3 - 1)}{n_2 + 1} f(n_1, n_2 + 2, n_3 - 2; 0, 0, 0) \\
& + (n_1 + n_2 + 2 n_3 + 2) f(n_1, n_2, n_3; 0, 0, 0) \\
& + \frac{1}{n_2 + 1} f(n_1, 2 + n_2, n_3; \star, 0, 0) \\
& + \frac{n_3 (n_3 - 1)}{n_1 + 1} f(n_1 + 2, n_2, n_3 - 2; 0, 0, 0) \\
& + \frac{1}{n_1 + 1} f(n_1 + 2, n_2, n_3; 0, \star, 0) \\
& - \frac{\delta_{n_3}}{n_2 + 1} f(n_1, n_2 + 2, \star; 0, 0, 0) \\
& - \frac{\delta_{n_3}}{n_1 + 1} f(n_1 + 2, n_2, \star; 0, 0, 0) \Big] \Big\}. \tag{4}
\end{aligned}$$

Formulas for  $f(n_1, n_2 + 2, n_3, 0, 0, 0)$  and  $f(n_1 + 2, n_2, n_3, 0, 0, 0)$  can be obtained from Eqs. (3) and (4) by interchanging  $(r_2, r_3)$  or  $(r_1, r_3)$  respectively.

The next three recursions increase by 1 any of  $n_4, n_5, n_6$  of a given subshell  $n_4 + n_5 + n_6$  for arbitrary  $n_1, n_2$  and  $n_3$

$$\begin{aligned}
f(n_1, n_2, n_3; n_4, n_5, n_6 + 1) = & \frac{1}{w_1 w_2 w_3} \{ \\
& (n_1 - 1) n_1 n_4 f(n_1 - 2, n_2, n_3; n_4 - 1, n_5 + 1, n_6) \\
& + (n_2 - 1) n_2 n_5 f(n_1, n_2 - 2, n_3; n_4 + 1, n_5 - 1, n_6) \\
& - (n_3 - 1) n_3 n_4 f(n_1, n_2, n_3 - 2; n_4 - 1, n_5 + 1, n_6) \\
& - (n_3 - 1) n_3 n_5 f(n_1, n_2, n_3 - 2; n_4 + 1, n_5 - 1, n_6) \\
& + n_4 n_5 (n_1 + n_2 - n_3 + n_6 + 1) f(n_1, n_2, n_3; n_4 - 1, n_5 - 1, n_6) \\
& - (n_1 - 1) n_1 f(n_1 - 2, n_2, n_3; n_4, n_5 + 1, n_6) w_1 \\
& + (n_3 - 1) n_3 f(n_1, n_2, n_3 - 2; n_4, n_5 + 1, n_6) w_1 \\
& - n_5 (n_1 + n_2 - n_3 + n_6 + 1) f(n_1, n_2, n_3; n_4, n_5 - 1, n_6) w_1 \\
& - (n_2 - 1) n_2 f(n_1, n_2 - 2, n_3; n_4 + 1, n_5, n_6) w_2 \\
& + (n_3 - 1) n_3 f(n_1, n_2, n_3 - 2; n_4 + 1, n_5, n_6) w_2 \\
& - n_4 (n_1 + n_2 - n_3 + n_6 + 1) f(n_1, n_2, n_3; n_4 - 1, n_5, n_6) w_2
\end{aligned}$$

$$\begin{aligned}
& -n_4 n_5 f(n_1, n_2, n_3; n_4 - 1, n_5 - 1, n_6 + 1) w_3 \\
& + (n_1 + n_2 - n_3 + n_6 + 1) f(n_1, n_2, n_3; n_4, n_5, n_6) w_1 w_2 \\
& + n_4 f(n_1, n_2, n_3; n_4 - 1, n_5, n_6 + 1) w_2 w_3 \\
& + n_5 f(n_1, n_2, n_3; n_4, n_5 - 1, n_6 + 1) w_3 w_1 \\
& - \delta_{n_1} n_4 f(\star, n_2, n_3; n_4 - 1, n_5 + 1, n_6) \\
& + \delta_{n_1} f(\star, n_2, n_3; n_4, n_5 + 1, n_6) w_1 \\
& - \delta_{n_2} n_5 f(n_1, \star, n_3; n_4 + 1, n_5 - 1, n_6) \\
& + \delta_{n_2} f(n_1, \star, n_3; n_4 + 1, n_5, n_6) w_2 \\
& + \delta_{n_3} n_4 f(n_1, n_2, \star; n_4 - 1, n_5 + 1, n_6) \\
& + \delta_{n_3} n_5 f(n_1, n_2, \star; n_4 + 1, n_5 - 1, n_6) \\
& - \delta_{n_3} f(n_1, n_2, \star; n_4, n_5 + 1, n_6) w_1 \\
& - \delta_{n_3} f(n_1, n_2, \star; n_4 + 1, n_5, n_6) w_2 \}.
\end{aligned} \tag{5}$$

Two other recursions for  $f(n_1, n_2, n_3, n_4, n_5 + 1, n_6)$  and  $f(n_1, n_2, n_3, n_4 + 1, n_5, n_6)$  are obtained from Eqs. (3) and (5) by interchanging arguments of  $f$ -function, using the identities

$$\begin{aligned}
& f(n_1, n_2, n_3; n_4, n_5, n_6; w_1, w_2, w_3) = \\
& f(n_2, n_1, n_3; n_5, n_4, n_6; w_2, w_1, w_3) = \\
& f(n_3, n_2, n_1; n_6, n_5, n_4; w_3, w_2, w_1) = \\
& f(n_1, n_3, n_2; n_4, n_6, n_5; w_1, w_3, w_2),
\end{aligned} \tag{6}$$

where any of  $n_i$  can become a  $\star$ .

### III. BOUNDARY TERMS

These recursion relations in Eqs. (4) and (5) involve boundary terms. The first boundary term was derived in Ref. [9], the 7 remaining boundary terms to start the first set of recursions are obtained from the generic Fromm-Hill formula [5] or by direct integration

$$\begin{aligned}
f(0, 0, 0; 0, 0, 0) = & \frac{-1}{2 w_1 w_2 w_3} \left\{ \right. \\
& \ln\left(\frac{w_3}{w_1 + w_2}\right) \ln\left(1 + \frac{w_3}{w_1 + w_2}\right) + \text{Li}_2\left(-\frac{w_3}{w_1 + w_2}\right) + \text{Li}_2\left(1 - \frac{w_3}{w_1 + w_2}\right)
\end{aligned}$$

$$\begin{aligned}
& + \ln\left(\frac{w_2}{w_3 + w_1}\right) \ln\left(1 + \frac{w_2}{w_1 + w_3}\right) + \text{Li}_2\left(-\frac{w_2}{w_1 + w_3}\right) + \text{Li}_2\left(1 - \frac{w_2}{w_1 + w_3}\right) \\
& + \ln\left(\frac{w_1}{w_2 + w_3}\right) \ln\left(1 + \frac{w_1}{w_1 + w_2}\right) + \text{Li}_2\left(-\frac{w_1}{w_2 + w_3}\right) + \text{Li}_2\left(1 - \frac{w_1}{w_2 + w_3}\right) \Big\}, \\
f(1, 0, 0; 0, 0, 0) &= -\frac{1}{w_2^2 w_3^2} \ln\left[\frac{w_1 (w_1 + w_2 + w_3)}{(w_1 + w_2)(w_1 + w_3)}\right], \\
f(0, 1, 0; 0, 0, 0) &= -\frac{1}{w_1^2 w_3^2} \ln\left[\frac{w_2 (w_1 + w_2 + w_3)}{(w_2 + w_3)(w_2 + w_1)}\right], \\
f(0, 0, 1; 0, 0, 0) &= -\frac{1}{w_1^2 w_2^2} \ln\left[\frac{w_3 (w_1 + w_2 + w_3)}{(w_3 + w_1)(w_3 + w_2)}\right], \\
f(1, 1, 0; 0, 0, 0) &= \frac{1}{w_1 w_2 (w_1 + w_2) w_3^2}, \\
f(1, 0, 1; 0, 0, 0) &= \frac{1}{w_1 w_3 (w_1 + w_3) w_2^2}, \\
f(0, 1, 1; 0, 0, 0) &= \frac{1}{w_2 w_3 (w_2 + w_3) w_1^2}, \\
f(1, 1, 1; 0, 0, 0) &= \frac{1}{w_1^2 w_2^2 w_3^2}, \tag{7}
\end{aligned}$$

where  $\text{Li}_2(x)$  is the dilogarithmic function. Besides the initial terms one needs also as boundary terms the following two-electron integrals

$$\begin{aligned}
f(\star, n_2, n_3; n_4, n_5, n_6) &= \Gamma(n_5 + n_6 - 1, n_4, n_3 + n_2 - 1; w_2 + w_3, w_1, 0), \\
f(n_1, \star, n_3; n_4, n_5, n_6) &= \Gamma(n_4 + n_6 - 1, n_5, n_1 + n_3 - 1; w_1 + w_3, w_2, 0), \\
f(n_1, n_2, \star; n_4, n_5, n_6) &= \Gamma(n_4 + n_5 - 1, n_6, n_1 + n_2 - 1; w_1 + w_2, w_3, 0), \\
f(n_1, n_2, n_3; \star, 0, 0) &= \Gamma(n_3 - 1, n_2 - 1, n_1; w_2, w_3, 0), \\
f(n_1, n_2, n_3; 0, \star, 0) &= \Gamma(n_1 - 1, n_3 - 1, n_2; w_3, w_1, 0), \\
f(n_1, n_2, n_3; 0, 0, \star) &= \Gamma(n_2 - 1, n_1 - 1, n_3; w_1, w_2, 0), \tag{8}
\end{aligned}$$

where  $\Gamma$  is defined in Eq. (A5). Recursion relations for  $\Gamma$  have already been worked out in the literature and can be found for example in [4, 10]. For completeness we include below generic formulas for two cases: nonnegative  $n_1, n_2, n_3$ ,

$$\Gamma(n_1, n_2, n_3; \alpha_1, \alpha_2, \alpha_3) = \left(-\frac{d}{d\alpha_1}\right)^{n_1} \left(-\frac{d}{d\alpha_2}\right)^{n_2} \left(-\frac{d}{d\alpha_3}\right)^{n_3} \frac{1}{(\alpha_1 + \alpha_2)(\alpha_2 + \alpha_3)(\alpha_3 + \alpha_1)}, \tag{9}$$

and the second case:  $n_1 = -1$ , and nonnegative  $n_2, n_3$

$$\Gamma(-1, n_2, n_3; \alpha_1, \alpha_2, \alpha_3) = \left(-\frac{d}{d\alpha_2}\right)^{n_2} \left(-\frac{d}{d\alpha_3}\right)^{n_3} \frac{\ln(\alpha_1 + \alpha_2) - \ln(\alpha_1 + \alpha_3)}{(\alpha_2 - \alpha_3)(\alpha_2 + \alpha_3)}. \tag{10}$$

The complete set of recursion relations have been checked by comparison with analytic expression obtained by differentiation of Fromm and Hill formula [5] with respect to  $u_i$  and  $w_i$  parameters (in our notation).

#### IV. DERIVATION OF THREE-ELECTRON RECURSION FORMULAS

We use the method of the integration by parts identities [11], which is by now standard in the analytical calculation of Feynman diagrams. In our case, it amounts to consider the following 9 identities in the momentum representation of the integral  $G$ , [see Eq. (A6)]

$$0 \equiv \text{id}(i, j) = \int d^3 k_1 \int d^3 k_2 \int d^3 k_3 \frac{\partial}{\partial \vec{k}_i} \left[ \vec{k}_j (k_1^2 + u_1^2)^{-m_1} (k_2^2 + u_2^2)^{-m_2} (k_3^2 + u_3^2)^{-m_3} (k_{32}^2 + w_1^2)^{-m_4} (k_{13}^2 + w_2^2)^{-m_5} (k_{21}^2 + w_3^2)^{-m_6} \right], \quad (11)$$

which are trivially valid (the integral of the derivative of a function vanishing at infinity vanishes).

These identities group naturally into three sets. The first set consists of  $\text{id}(1, 1)$ ,  $\text{id}(2, 1)$ , and  $\text{id}(3, 1)$ . Other sets are obtained by changing the second argument from 1 into 2 or 3. The reduction of the scalar products from numerator leads to the following identities of the first set

$$\begin{aligned} \text{id}(1, 1) = & -m_6 G(m_1 - 1, m_6 + 1) - m_5 G(m_1 - 1, m_5 + 1) + m_6 G(m_2 - 1, m_6 + 1) \\ & + m_5 G(m_3 - 1, m_5 + 1) + (3 - 2m_1 - m_5 - m_6) G() + m_6 G(m_6 + 1) u_1^2 \\ & + m_5 G(m_5 + 1) u_1^2 + 2m_1 G(m_1 + 1) u_1^2 - m_6 G(m_6 + 1) u_2^2 - m_5 G(m_5 + 1) u_3^2 \\ & + m_5 G(m_5 + 1) w_2^2 + m_6 G(m_6 + 1) w_3^2, \end{aligned}$$

$$\begin{aligned} \text{id}(2, 1) = & -m_6 G(m_1 - 1, m_6 + 1) - m_5 G(m_1 - 1, m_5 + 1) + m_6 G(m_2 - 1, m_6 + 1) \\ & + m_5 G(m_3 - 1, m_5 + 1) - m_5 G(m_4 - 1, m_5 + 1) + (m_6 - m_1) G() \\ & + m_5 G(m_5 + 1, m_6 - 1) - m_1 G(m_1 + 1, m_2 - 1) + m_1 G(m_1 + 1, m_6 - 1) \\ & + m_6 G(m_6 + 1) u_1^2 + m_5 G(m_5 + 1) u_1^2 + m_1 G(m_1 + 1) u_1^2 - m_6 G(m_6 + 1) u_2^2 \\ & + m_1 G(m_1 + 1) u_2^2 - m_5 G(m_5 + 1) u_3^2 + m_5 G(m_5 + 1) w_1^2 - m_6 G(m_6 + 1) w_3^2 \\ & - m_5 G(m_5 + 1) w_3^2 - m_1 G(m_1 + 1) w_3^2, \end{aligned}$$

$$\begin{aligned}
\text{id}(3, 1) = & -m_6 G(m_1 - 1, m_6 + 1) - m_5 G(m_1 - 1, m_5 + 1) + m_6 G(m_2 - 1, m_6 + 1) \\
& + m_5 G(m_3 - 1, m_5 + 1) - m_6 G(m_4 - 1, m_6 + 1) + m_6 G(-m_5 + 1, m_6 + 1) \\
& + (m_5 - m_1) G() - m_1 G(m_1 + 1, m_3 - 1) + m_1 G(m_1 + 1, m_5 - 1) \\
& + m_6 G(m_6 + 1) u_1^2 + m_5 G(m_5 + 1) u_1^2 + m_1 G(m_1 + 1) u_1^2 - m_6 G(m_6 + 1) u_2^2 \\
& - m_5 G(m_5 + 1) u_3^2 + m_1 G(m_1 + 1) u_3^2 + m_6 G(m_6 + 1) w_1^2 - m_6 G(m_6 + 1) w_2^2 \\
& - m_5 G(m_5 + 1) w_2^2 - m_1 G(m_1 + 1) w_2^2.
\end{aligned} \tag{12}$$

The function  $G$  is defined as in Eq. (A6), but for ease of writing only the arguments which contain a  $\pm 1$  are shown explicitly [so that, for instance,  $G(m_5 + 1)$  stands for  $G(m_1, m_2, m_3; m_4, m_5 + 1, m_6)$  etc.]

The general solution of these recursions is on itself of great interest, but we consider here only the case  $u_1 = u_2 = u_3 = 0$  which corresponds to Hylleraas basis set. For this we put  $m_1 = m_2 = m_3 = 1$  and differentiate these identities over  $u_1, u_2, u_3$  at  $u_1 = u_2 = u_3 = 0$ . It leads to recursions for the  $h$  function of Eq. (A7). The first subset of identities forms 3 linear equations for  $h(n_1 + 2), h(m_5 + 1), h(m_6 + 1)$  (same convention for the explicitly written indices as for the function  $G$ ) which can easily be solved. Since there are 9 equations for 6 unknowns, two different solutions for  $h(m_4 + 1), h(m_5 + 1), h(m_6 + 1)$  can be used to simplify recursion formulas. We achieve this by solving three equations

$$\begin{aligned}
h_1(m_4 + 1) - h_2(m_4 + 1) &= 0, \\
h_1(m_5 + 1) - h_2(m_5 + 1) &= 0, \\
h_1(m_6 + 1) - h_2(m_6 + 1) &= 0,
\end{aligned} \tag{13}$$

against the following linear combinations

$$\begin{aligned}
& h(m_5 - 1, m_6 + 1) - h(m_4 - 1, m_6 + 1), \\
& h(m_5 + 1, m_6 - 1) - h(m_4 - 1, m_5 + 1), \\
& h(m_4 + 1, m_5 - 1) - h(m_4 + 1, m_6 - 1).
\end{aligned} \tag{14}$$

The obtained solutions are inserted back in recursion formulas for  $h(n_i + 2)$  and  $h(m_i + 1)$ , and they take the form

$$\frac{2 w_2^2 w_3^2}{n_1 + 1} h(n_1 + 2) = [w_3^2 (2 + n_1 + n_2 + 2 n_3) + w_2^2 (2 + n_1 + 2 n_2 + n_3) -$$



$$\begin{aligned}
& w_1^2 (2 + 2n_1 + n_2 + n_3)] h() + w_3^2 Q_3 + w_2^2 Q_2 - w_1^2 Q_1 + \dots \\
\frac{2w_1^2 w_3^2}{n_2 + 1} h(n_2 + 2) &= [w_3^2 (2 + n_1 + n_2 + 2n_3) - w_2^2 (2 + n_1 + 2n_2 + n_3) + \\
& w_1^2 (2 + 2n_1 + n_2 + n_3)] h() + w_3^2 Q_3 - w_2^2 Q_2 + w_1^2 Q_1 + \dots \\
\frac{2w_1^2 w_2^2}{n_3 + 1} h(n_3 + 2) &= [-w_3^2 (2 + n_1 + n_2 + 2n_3) + w_2^2 (2 + n_1 + 2n_2 + n_3) + \\
& w_1^2 (2 + 2n_1 + n_2 + n_3)] h() - w_3^2 Q_3 + w_2^2 Q_2 + w_1^2 Q_1 + \dots \quad (15)
\end{aligned}$$

$$\begin{aligned}
w_1^2 m_4 h(m_4 + 1) &= (-n_1 + n_2 + n_3 + 2m_4 - 1) h() + 2(X_3 - X_2) + \dots \\
w_2^2 m_5 h(m_5 + 1) &= (n_1 - n_2 + n_3 + 2m_5 - 1) h() + 2(X_1 - X_3) + \dots \\
w_3^2 m_6 h(m_6 + 1) &= (n_1 + n_2 - n_3 + 2m_6 - 1) h() + 2(X_2 - X_1) + \dots \quad (16)
\end{aligned}$$

where by  $\dots$  we denote the omitted boundary terms, which are proportional to  $\delta_{n_i}$ , and

$$\begin{aligned}
Q_1 &= \frac{n_1(n_1 - 1)}{n_3 + 1} h(n_1 - 2, n_3 + 2) + \frac{n_1(n_1 - 1)}{n_2 + 1} h(n_1 - 2, n_2 + 2) + \\
& \frac{1}{n_3 + 1} h(n_3 + 2, m_5 - 1) + \frac{1}{n_2 + 1} h(n_2 + 2, m_6 - 1), \\
Q_2 &= \frac{n_2(n_2 - 1)}{n_3 + 1} h(n_2 - 2, n_3 + 2) + \frac{n_2(n_2 - 1)}{n_1 + 1} h(n_1 + 2, n_2 - 2) + \\
& \frac{1}{n_3 + 1} h(n_3 + 2, m_4 - 1) + \frac{1}{n_2 + 1} h(n_1 + 2, m_6 - 1), \\
Q_3 &= \frac{n_3(n_3 - 1)}{n_2 + 1} h(n_2 + 2, n_3 - 2) + \frac{n_3(n_3 - 1)}{n_1 + 1} h(n_1 + 2, n_3 - 2) + \\
& \frac{1}{n_2 + 1} h(n_2 + 2, m_4 - 1) + \frac{1}{n_1 + 1} h(n_1 + 2, m_5 - 1), \quad (17)
\end{aligned}$$

$$\begin{aligned}
X_1 &= (n_2 - 1) n_2 m_4 h(n_2 - 2, m_4 + 1) - (n_3 - 1) n_3 m_4 h(n_3 - 2, m_4 + 1), \\
X_2 &= (n_3 - 1) n_3 m_5 h(n_3 - 2, m_5 + 1) - (n_1 - 1) n_1 m_5 h(n_1 - 2, m_5 + 1), \\
X_3 &= (n_1 - 1) n_1 m_6 h(n_1 - 2, m_6 + 1) - (n_2 - 1) n_2 m_6 h(n_2 - 2, m_6 + 1). \quad (18)
\end{aligned}$$

There is obviously some freedom in using Eqs. (13) to simplify recursions, and our choice allows to separate recursions into two subshells of  $n_1 + n_2 + n_3$  and  $m_4 + m_5 + m_6$ . We put  $m_4 = m_5 = m_6 = 1$  in Eq. (15) and immediately obtain the first subset of recursions, Eq. (4). The second subset Eq. (5) is obtained from Eq. (16) in three steps. The first step is setting  $m_4 = m_5 = m_6 = 1$ . The second step is multiplication of recursion formulas by  $w_i w_j$  to remove any  $w_i$  from the denominator. The third step is differentiation with respect to  $w_1, w_2$ , and  $w_3$ , which converts the function  $h()$  into the function  $f()$ . That completes the derivation of recursion relations for three-electron Hylleraas integrals.

## V. SUMMARY

We have derived a complete set of recursion relations for three-electron Hylleraas integral  $f$ . They are sufficiently stable for a precise numerical calculation. Since all  $w_i$  are bounded from below by approximately  $\sqrt{2E}$ , where  $E$  is the ionization energy, the denominators of three-electron recursions are also bounded from below. Therefore, by using extended precision arithmetics (octuple precision) we can safely devote 1 digit per iteration, and still preserve quad precision for about 30 iterations, which is the maximum we aim to use. Possible instabilities in two-electron integrals are avoided by applying inverse recursions [10]. Numerical results are shown in Table I. These recursion relations can be used for high

TABLE I: Values of three-electron Hylleraas integral at  $w_1 = w_2 = w_3 = 1$ ,  $[m]$  denotes  $10^m$

| $n$ | $f(n, 0, 0; 0, 0, 0)$                 | $f(0, 0, 0; n, 0, 0)$                 |
|-----|---------------------------------------|---------------------------------------|
| 0   | 2.208 310 154 388 618 874 536 424[-1] | 2.208 310 154 388 618 874 536 424[-1] |
| 1   | 2.876 820 724 517 809 274 392 190[-1] | 2.208 310 154 388 618 874 536 424[-1] |
| 2   | 6.071 253 765 587 525 062 881 067[-1] | 3.658 582 716 243 175 207 969 277[-1] |
| 3   | 1.801 456 579 614 247 419 513 752[0]  | 8.803 723 087 040 150 596 505 449[-1] |
| 4   | 6.949 688 537 201 117 333 162 822[0]  | 2.849 464 173 126 685 211 199 798[0]  |
| 5   | 3.316 893 553 498 521 645 367 878[1]  | 1.176 795 411 671 425 935 279 582[1]  |
| 6   | 1.892 427 697 247 010 803 401 964[2]  | 5.962 899 567 501 152 778 486 008[1]  |
| 7   | 1.258 719 915 821 483 876 136 660[3]  | 3.596 955 116 745 326 378 301 909[2]  |
| 8   | 9.575 385 319 725 442 534 735 866[3]  | 2.522 862 411 307 814 783 043 058[3]  |
| 9   | 8.206 804 555 680 135 296 239 238[4]  | 2.019 476 554 953 447 619 512 494[4]  |

precision calculation of lithium wave function with a small computational effort. We think that similar recursions can be derived also for beryllium, which involves four electrons, but the generic integral with all powers of  $r_{ij}$  equal to  $-1$  has not yet been obtained. However, this generic integral can be represented as a multiple sum as recently indicated by Sims and Hagstrom [12] and Frolov [13] and thus obtained numerically. Coming back to lithium, our aim is the high precision calculation of higher order relativistic and QED effects like that for lithium hyperfine splitting. It requires, however, also the calculation of Hylleraas integrals with various negative powers of  $r_i$  and  $r_{ij}$ .  $1/r_i^n$  with  $n > 1$  can be expressed in terms of

harmonic polylogarithms, which were introduced recently in [14], and  $1/r_{ij}^n$  can be obtained, we think, by generalization of the recursion relations obtained in this work.

In summary, we have presented a simple method to derive recursion relations and solved them for three-electron Hylleraas integral.

## VI. ACKNOWLEDGMENTS

We are grateful to Vladimir Korobov for his source code of the fast multiprecision arithmetics and of two-electron recursions in the singular case. This work was supported by EU grant HPRI-CT-2001-50034.

- 
- [1] E.A. Hylleraas, Z. Phys. **54**, 347 (1929).
  - [2] G.W.F. Drake, in *Long Range Casimir Forces: Theory and Recent Experiments on Atomic Systems*, ed. F.S. Levin and D.A. Micha, Plenum, New York, 1993 p. 107.
  - [3] G.W.F. Drake and Z.-C. Yan, Phys. Rev. A **46**, 2378 (1992); Z.-C. Yan and G. W. F. Drake, Phys. Rev. A **52**, 3711 (1995); Phys. Rev. A **61**, 022504 (2000); Phys. Rev. A **66**, 042504 (2002); Phys. Rev. Lett. **91**, 113004 (2003).
  - [4] R.A. Sack, C.C.J Roothaan and W. Kołos, J. Math. Phys. **8**, 1093 (1967).
  - [5] D. M. Fromm and R. N. Hill, Phys. Rev. A **36**, 1013 (1987).
  - [6] F. E. Harris, Phys. Rev. A **55**, 1820 (1997).
  - [7] V.S. Zotev and T.K. Rebane. Phy. Rev. A **65**, 062501 (2002).
  - [8] F.E. Harris, A.M. Frolov, and V.H. Smith, Jr., J. Chem. Phys. **119**, 8833 (2003).
  - [9] E. Remiddi, Phys. Rev. A **44**, 5492 (1991).
  - [10] V.I. Korobov, J. Phys. B **35**, 1959 (2002).
  - [11] F.V. Tkachov, Phys. Lett. **B100**, 65 (1981); K.G. Chetyrkin and F.V. Tkachov, Nucl. Phys. B19281159.
  - [12] J.S. Sims and S.A. Hagstrom, J. Phys. B **37**, 1519 (2004).
  - [13] A.M. Frolov, J. Phys. B. **37**, 2103 (2004).
  - [14] E. Remiddi and J.A.M. Vermaseren, Int. J. Mod. Phys. **A15**, 7275 (2000).

## APPENDIX A: DEFINITIONS OF FUNCTIONS AND RELATIONS

We use the symbol  $\delta_n$  which denotes the Kronecker  $\delta_{n,0}$ . Definitions of all functions used in this work are presented below.  $r_{ij}$  denotes  $|\vec{r}_i - \vec{r}_j|$  and  $k_{ij} = |\vec{k}_i - \vec{k}_j|$ . All the  $n_i$  and  $m_i$  are assumed to be nonnegative integer and  $w_i, u_i$  to be real and positive.

$$F(n_1, n_2, n_3; n_4, n_5, n_6) = \int \frac{d^3 r_1}{4\pi} \int \frac{d^3 r_2}{4\pi} \int \frac{d^3 r_3}{4\pi} e^{-w_1 r_1 - w_2 r_2 - w_3 r_3 - u_1 r_{23} - u_2 r_{31} - u_3 r_{12}} r_{23}^{n_1-1} r_{31}^{n_2-1} r_{12}^{n_3-1} r_1^{n_4-1} r_2^{n_5-1} r_3^{n_6-1} \quad (\text{A1})$$

$$f(n_1, n_2, n_3; n_4, n_5, n_6) = \int \frac{d^3 r_1}{4\pi} \int \frac{d^3 r_2}{4\pi} \int \frac{d^3 r_3}{4\pi} e^{-w_1 r_1 - w_2 r_2 - w_3 r_3} r_{23}^{n_1-1} r_{31}^{n_2-1} r_{12}^{n_3-1} r_1^{n_4-1} r_2^{n_5-1} r_3^{n_6-1} \quad (\text{A2})$$

$$f(\star, n_2, n_3; n_4, n_5, n_6) = \int \frac{d^3 r_1}{4\pi} \int \frac{d^3 r_2}{4\pi} \int \frac{d^3 r_3}{4\pi} e^{-w_1 r_1 - w_2 r_2 - w_3 r_3} 4\pi \delta^3(r_{23}) r_{31}^{n_2-1} r_{12}^{n_3-1} r_1^{n_4-1} r_2^{n_5-1} r_3^{n_6-1} \quad (\text{A3})$$

$$f(n_1, n_2, n_3; \star, n_5, n_6) = \int \frac{d^3 r_1}{4\pi} \int \frac{d^3 r_2}{4\pi} \int \frac{d^3 r_3}{4\pi} e^{-w_1 r_1 - w_2 r_2 - w_3 r_3} r_{23}^{n_1-1} r_{31}^{n_2-1} r_{12}^{n_3-1} 4\pi \delta^3(r_1) r_2^{n_5-1} r_3^{n_6-1} \quad (\text{A4})$$

$$\Gamma(n_1, n_2, n_3; \alpha_1, \alpha_2, \alpha_3) = \int \frac{d^3 r_1}{4\pi} \int \frac{d^3 r_2}{4\pi} e^{-\alpha_1 r_1 - \alpha_2 r_2 - \alpha_3 r_{12}} r_1^{n_1-1} r_2^{n_2-1} r_{12}^{n_3-1} \quad (\text{A5})$$

$$G(m_1, m_2, m_3; m_4, m_5, m_6) = \frac{1}{8\pi^6} \int d^3 k_1 \int d^3 k_2 \int d^3 k_3 (k_1^2 + u_1^2)^{-m_1} (k_2^2 + u_2^2)^{-m_2} (k_3^2 + u_3^2)^{-m_3} (k_{32}^2 + w_1^2)^{-m_4} (k_{13}^2 + w_2^2)^{-m_5} (k_{21}^2 + w_3^2)^{-m_6} \quad (\text{A6})$$

$$h(n_1, n_2, n_3; m_4, m_5, m_6) = (-1)^{n_1} \frac{\partial^{n_1}}{\partial u_1^{n_1}} \Big|_{u_1=0} (-1)^{n_2} \frac{\partial^{n_2}}{\partial u_2^{n_2}} \Big|_{u_2=0} (-1)^{n_3} \frac{\partial^{n_3}}{\partial u_3^{n_3}} \Big|_{u_3=0} G(1, 1, 1, m_4, m_5, m_6) \quad (\text{A7})$$

$$f(n_1, n_2, n_3; n_4, n_5, n_6) = (-1)^{n_4} \frac{\partial^{n_4}}{\partial w_1^{n_4}} (-1)^{n_5} \frac{\partial^{n_5}}{\partial w_2^{n_5}} (-1)^{n_6} \frac{\partial^{n_6}}{\partial w_3^{n_6}} h(n_1, n_2, n_3, 1, 1, 1) \quad (\text{A8})$$